

Transtect™

Active Ingredient:	By Wt
*Dinotefuran.....	70%
Other Ingredients:	30%
Total:	100%

*N-methyl-N'-nitro-N"-[(tetrahydro-3-furanyl)methyl]guanidine

KEEP OUT OF REACH OF CHILDREN CAUTION

For foliar and systemic control of sucking and chewing insects infesting ornamental plants grown in outdoor landscapes (commercial, industrial, recreational and residential), parks, schools, cemeteries, airport grounds, military grounds, golf courses, woodlots, athletic fields, forests (National, Private and State) and other wooded areas.

For outdoor use only

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

EPA Reg. No. 59639-170-74779

EPA Est. 46275-IL-2

NET WEIGHT: 0.75 POUNDS (CONTAINS 20/0.6 OZ WSP PACKETS)

Manufactured for:



Rainbow Treecare Scientific Advancements

11571 K-Tel Dr.

Minnetonka, MN 55343

1-877-ARBORIST

1-877-272-6747

www.treecarescience.com

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> Call poison control center or doctor immediately for treatment advice. Do not induce vomiting unless told to by the poison control center or doctor. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.
IF INHALED:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call poison control center or doctor for further treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	
<p>HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal), or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident). Call CHEMTREC at 1-800-424-9300.</p>	

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

USER SAFETY REQUIREMENTS:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not dispose of equipment washwaters or rinsate into a natural drain or water body. Do not contaminate water when disposing of equipment washwaters or rinsate.

This compound is toxic to honey bees. The persistence of residues and potential residual toxicity of Transtect 70WSP in nectar and pollen suggest the possibility of chronic risk to honey bee larvae and the eventual instability of the hive.

- This product is toxic to bees exposed to residues for more than 38 hours following treatment.
- Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state and federal authorities.

Transtect 70WSP and its degradate, MNG, have the properties and characteristics associated with chemicals detected in groundwater. The high water solubility of Transtect 70WSP, and its degradate, MNG, coupled with its very high mobility, and resistance to biodegradation indicates that this compound has a strong potential to leach to the subsurface under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Periodic monitoring of shallow groundwater in the use area is recommended.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

SPRAY DRIFT ADVISORY

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crop thereof rendered for sale, use or consumption.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL, USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

BEE HAZARD DIRECTIONS FOR USE

Follow these additional application restrictions for use patterns marked with the Bee Hazard Icon.



1. FOR COMMERCIALLY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS

Do not apply this product while bees are foraging.

- This product is toxic to bees exposed to residues for more than 38 hours following treatment.
- Do not apply this product to blooming, pollen-shedding or nectar producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state or federal authorities.

Follow these additional application restrictions for use patterns marked with the Bee Hazard Icon.



2. NON-AGRICULTURAL PRODUCTS:

Do not apply Transtect 70WSP while bees are foraging. Do not apply Transtect 70WSP to plants that are flowering. Only apply after all flower petals have fallen off.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.

Look for the bee hazard icon  in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>. Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried. Keep all unprotected persons out of operating areas, or vicinity where there may be drift.

**DISCLAIMER, RISKS OF USING THIS PRODUCT, LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and subject to the Risks of Using This Product as described above. **To the extent consistent with applicable law AND AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

Valent must be provided with prompt notice of any claim so that an immediate inspection of the affected property or plants can be made. To the extent applicable law allows such requirement, notice of a claim shall be given promptly to Valent, whether the claim is based on contract, negligence, strict liability, or other tort or otherwise be barred from any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.

RESISTANCE MANAGEMENT RECOMMENDATIONS

Transtect 70WSP contains a Group 4A insecticide. Insect biotypes with acquired resistance to Group 4A may eventually dominate the insect population if Group 4A insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Transtect 70WSP or other Group 4A insecticides.

To delay insecticide resistance consider:

- Avoiding the consecutive use of Dinotefuran Tree Care 70WSP, or other Group 4A insecticides that have similar target site of action, on the same insect species.
- Using tank mixtures or premixes with insecticides from a different tank target site of action Group as long as the involved products are all registered for the same use and have different sites of action.
- Basing insecticide use on a comprehensive IPM program.
- Monitoring treated insect populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturers for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.

For further information, contact Valent U.S.A. Corporation at the following toll free number: 1-800-682-5368

APPLICATION INFORMATION

Transtect 70WSP is a systemic insecticide that is taken up by the root system, foliage or through the bark and translocated upward throughout the plant. Apply Transtect 70WSP by soil injection, soil drench, basal trunk spray or foliar spray to ornamental plants growing in outdoor landscapes (commercial, industrial, recreational, and residential), parks, schools, cemeteries, airport grounds, military grounds, golf courses, woodlots, athletic fields, forests (National, Private and State) and other wooded areas.

Note: The active ingredient in Transtect 70WSP has been found to be effective on a wide range of insect species. Before using Transtect 70WSP for insects not listed in the directions for use, test Transtect 70WSP on a small scale basis and evaluate for insect control prior to widespread use.

USE PRECAUTIONS

- Do not apply when the target site is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist.

RESTRICTIONS

- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Do not allow runoff or puddling of irrigation water following application.
- Do not apply to areas which are water logged or saturated or frozen which will not allow penetration into the root zone of the plant.
- When making foliar applications or basal trunk sprays keep children and pets away from the treated area until the spray has dried.
- Do not apply this product, by any application method, to linden, basswood or other Tilia species.

MIXING INSTRUCTIONS:

Transtect 70WSP Alone: Add 1/2 of the required amount of water to the mix tank. With the agitator running, add the desired amount of Transtect 70WSP to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after Transtect 70WSP has completely disappeared into the mix water. Maintain agitation until all of the mixture has been applied.

Transtect 70WSP + Tank Mixtures: Add 1/2 of the required amount of water to the mix tank. Start the agitator before adding any tank mix partners. Add tank mix partners in this order: products packaged in water soluble packaging, wettable powders, wettable granules (dry flowables), liquid flowables, liquids, emulsifiable concentrates, surfactants, and adjuvants. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all the mixture has been applied.

Note: When tank mixing, add the Transtect 70WSP water soluble packets first, followed by any additional products in water soluble packaging. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using Transtect 70WSP in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions and limitations which appear on the tank mix product label. Do not exceed the label dosage rate, and follow the most restrictive label precautions and limitations. Do not mix this product with any product which prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

COMPATIBILITY

IMPORTANT: The risk factors of all potential tank mixes have not been tested on all plants. Before applying any tank mixture not specifically recommended on this label, confirm risk factors to the target plant.

Transtect 70WSP is compatible with most commonly used pesticides, adjuvants and nutritional sprays. However, since it is not possible to test all possible mixtures, pre-test to assure the physical compatibility and lack of phytotoxic effect of any proposed mixture with Transtect 70WSP. To determine the physical compatibility of Transtect 70WSP with other products, use the test described below. When an adjuvant is to be used with this product, Valent recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

In an appropriately sized container, add 1 (0.6 oz) packet of Transtect 70WSP along with the proportionate amounts of additional products to 5 qt of water, add wettable powders and water dispersible granular products first, then liquid flowables and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for additional required ingredients to the spray tank.

APPLICATION TECHNIQUES

Notes:

- Use higher labeled rates for broadleaf evergreens with dense foliage (example hollies), for applications on large trees, for difficult to control insects, for therapeutic treatments, or for preventive applications made greater than 4 weeks in advance of the target pest infestation
- Transtect 70WSP water dilution rates range from 1 fl oz of water up to 1 gal of water per foot of shrub height or inch of trunk diameter. Specific dilution rates will depend further on the application equipment and use patterns for each individual.
- Keep soil moist for at least 7 days after application.
- Heavy rainfall or inadequate irrigation immediately following application will decrease performance.
- Applications made during severe drought conditions will result in slower uptake and translocation.

USE RESTRICTIONS

- Do not apply to excessively dry, saturated or frozen soil.

DETERMINING DIAMETER AT BREAST HEIGHT (DBH)

Single Stem: Measure the standard DBH of the tree at 4' 6" above the soil.

Multi Individual Trees Growing in Close Proximity: For trees that have grown close together, measure the DBH of each stem and treat each tree individually. Also, because of close proximity of trees, it may be necessary to apply Transtect 70WSP to the outer perimeter of clumped trees.

Multi-stem Split Below DBH: For a tree that has multiple stems splitting below DBH, measure the tree at the narrowest point between the root flare and the split.

Stem Clusters: For trees that are grown too close together to be treated as individual trees, measure the DBH of each stem and add the measurements together. Also, because of close proximity of trees, it may be necessary to apply Transtect 70WSP to the outer perimeter of clumped trees.

Tree Splits at DBH: For a tree that splits into two or more stems at DBH, measure and add the diameter of the stems and measure the narrowest point below the split. Take the average of these values.

SOIL APPLICATION

When applied to the soil, Transtect 70WSP is taken up by actively growing trees and shrubs. Speed of control will be dependent on tree size, tree health, environmental conditions and how actively pests are feeding. In actively growing and transpiring plants, control may be evident within one week after application in small shrubs, and within three weeks after application in large trees. For optimal control, time soil applications so that Transtect 70WSP has had time to move to insect feeding sites by time target life stage is present. Control may be less effective applied when soil is dry or when plants are not actively taking up water.

Remove landscape mulch, landscape fabric and surface organic matter prior to application to ensure solution is delivered to the mineral soil

Soil Drench: Measure the diameter of the tree. Mix required dose in water and uniformly apply to soil around the base of tree. Make applications within 12 inches of the trunk of the tree. Pull back mulch before drenching. For optimal performance, apply 1 – 4 pints of drench solution per foot of shrub height or inch of trunk diameter. Adjust drench volume based on soil type, soil moisture and thickness of mulch so that product is moved into root zone. Lower drench volumes may be less effective in dry soils or when applied over heavy mulch unless there is adequate rainfall or irrigation after application to move product into root zone.

Soil Injection: Measure the diameter of the tree. Mix required dose in water and apply to each shrub or tree using a soil injection application system. Transtect 70WSP may be soil injected with low volume or high volume injection equipment. For optimum performance, inject 1 to 32 fl oz of dilute solution per foot of shrub height or inch of trunk diameter depending on application equipment. Make shallow injections where feeder roots are most concentrated. Divide the required dose evenly among injection sites spaced as uniformly as possible around the tree. For trees less than 6 inches DBH, use at least 4 injection sites evenly spaced around the tree. Inject diluted solution approximately 2 to 6 inches deep into the mineral soil. Orient injection orifices to release the diluted product horizontally at the point of injection.

Basal System (preferred method): Apply Transtect 70WSP by placing the injections sites at the base of the tree. Evenly space the injection sites around the base of the tree. For most species a high density of fibrous roots are located at the base of the tree.

Grid System: Space injections on a 2.5 ft center within the drip line of the tree.

Circle System: Make injections every 2 to 3 feet within the dripline forming concentric circles around the base of the tree.

Applications should be made before trees are heavily infested and showing obvious symptoms of decline.

Note: Application to heavily infested trees may not prevent the decline or death of trees due to existing damage and tree stress.

Soil application for systemic insect control on ornamental plants in outdoor landscapes (commercial, industrial, recreational, and residential), parks, schools, cemeteries, airport grounds, military grounds, golf courses, woodlots, athletic fields, forests (National, Private and State) and other wooded areas.	
Pests	Rate
Adelgids (such as: Hemlock Woolly) Aphids Bagworms (larvae) Erythrina gall wasp Flatheaded borers (such as Emerald Ash Borer, Appletree) Glassy-Winged Sharpshooter Gypsy moth (larvae) Japanese beetle (adults) Lacebugs (such as: Azalea, Hawthorn) Leaf Beetles (such as Viburnum) Leafhoppers Leafminers (such as: Boxwood, Serpentine) Mealybugs (such as: Citrus, Long-tailed, Madeira, Obscure, Pink hibiscus, Root) Mimosa webworm (larvae) Peachtree borer (larvae) Pine tip moth (larvae) Plantbugs Psyllids (such as Asian citrus) Root Weevils (larvae and adults) (such as: Black vine) Roundheaded borers (excluding Asian Longhorned Beetle) Royal Palm Bug Sawfly larvae Scales (Armored and Soft) Spittlebug Tent caterpillars (larvae) Thrips (suppression) (such as: Chilli) Treehoppers Whiteflies (such as: Ficus, Giant, Silverleaf/Sweetpotato) White Grubs (such as: Oriental beetle)	<p>For Trees: Apply 1 (0.6 oz) packet of Dinotefuran Tree Care 70WSP per 5 to 17 inches of trunk diameter at breast height (DBH)</p> <p>For Shrubs: Apply 1 (0.6 oz) packet of Dinotefuran Tree Care 70WSP per 10 to 17 ft of height</p> <p>For Hedges: Apply 1 (0.6 oz) packet of Dinotefuran Tree Care 70WSP per foot of hedge height per 50 - 210 linear feet of row</p> <p>(Each packet contains approximately 0.026 lbs active ingredient)</p> <p>For Soil Drench: Combine Dinotefuran Tree Care 70WSP packets in sufficient water to achieve an application of at least 1 to 4 pints of dilute solution per foot of shrub height or inch of trunk diameter. For hedges, wet a one foot wide band down center of hedge row.</p> <p>For Soil Injection: Dinotefuran Tree Care 70WSP may be soil injected with low volume or high volume injection equipment. For optimum performance, inject 1 to 32 fl oz of dilute solution per foot of shrub height or inch of trunk diameter depending on application equipment. Make shallow injections where feeder roots are most concentrated.</p>
Do not apply more than 0.77 lbs (0.54 lbs ai) Transtect 70WSP per acre of landscape or forest per year.	



FOLIAR APPLICATION

Follow application instructions as indicated in the Bee Hazard Direction for Use

Apply in sufficient water to ensure good coverage of ornamental plants. In situations where a dense canopy exists and/or pest pressure is high, use greater water volumes. The use of a spray adjuvant may improve spray coverage. When making applications to plants with hard to wet foliage such as holly or pine, the addition of a spreader/sticker is recommended. Read and understand adjuvant product labels prior to mixing and applying them with Transtect 70WSP.

To assure optimum effectiveness, the product must be placed where the growing portion of the target plant can absorb the active ingredient. For optimal control, make application just before pest populations reach an economic threshold. If necessary, make a second application after 14 to 21 days.

Tank mixing with a surfactant may improve control of pests such as whitefly, mealybug, and scale. Confirm plant safety of tank mix in small area before using on a commercial scale.

Foliar application for systemic insect control on ornamental plants in outdoor landscapes (commercial, industrial, recreational and residential), parks, schools, cemeteries, airport grounds, military grounds, golf courses, woodlots, athletic fields, forests (National, Private and State) and other wooded areas.	
Pests	Rate
Adelgids (such as: Hemlock Woolly) Aphids (suppression) Glassy-Winged Sharpshooter Japanese beetle (adults) Lacebugs (such as: Azalea, Hawthorn) Leaf Beetles (such as Viburnum) Leafhoppers Leafminers (such as: Boxwood, Serpentine) Mealybugs (such as: Citrus, Long-tailed, Madeira, Obscure, Pink hibiscus, Root) Psyllids (such as Asian citrus) Root Weevils (adults) (such as: Black Vine) Scales (Armored and Soft) Thrips (suppression) (such as: Chilli) Whiteflies (such as: Ficus, Giant, Silverleaf/Sweetpotato)	Apply 1 (0.6 oz) packet of Dinotefuran Tree Care 70WSP per 25 to 50 gals (Each packet contains approximately 0.026 lbs active ingredient) 25 gallons of spray mix will treat approximately 5,000 square feet of area
Do not apply more than 0.77 lbs (0.54 lbs ai) Transtect 70WSP per acre of landscape or forest per year.	

BARK APPLICATION

When sprayed on the trunk, Transtect 70WSP will be absorbed through the bark and into the vascular system, and then transported throughout the tree. Speed of control will depend on tree size, tree health, environmental conditions and how actively pests are feeding. In actively transpiring trees, control may be evident within 1 – 4 weeks after application.

Spray bark on root flare (buttress roots) and on trunk between soil surface and 4 to 5 feet above the soil surface. Adjust nozzle to uniformly distribute spray over the entire circumference of the tree trunk and buttress roots. Wet bark just to the point of saturation and run off. Apply with a low volume sprayer operated at 10 to 20 PSI to prevent tree damage, bounce back and drift.

Application Equipment: Transtect 70WSP may be applied using many different types of low pressure sprayers. Before applying, start with a clean, well maintained sprayer. Apply a sufficient amount of water to ensure good coverage to the entire circumference of the lower 5 feet of the tree trunk. To reduce drift, use a spray nozzle that produces medium-sized droplets.

Mixing Instructions: Fill a clean spray tank to 2/3 of desired level with clean water. Add 6 – 12 packets (3.6 – 7.2 oz) of Transtect 70WSP per gallon of spray solution. A surfactant may be added to the spray solution to improve surface wetting and bark penetration. When using a surfactant, follow all label directions and precautions on the surfactant product label. Fill spray tank to desired level with clean water. Continue agitation until all spray solution has been applied. Prepare no more spray mixture than is required for immediate spray application.

Use a low pressure sprayer operated at 10 to 20 psi to reduce spray drift or splash back on non-target plants. Spray solution onto the tree trunk from 4" to 60" above the soil surface. Adjust sprayer nozzle to uniformly distribute spray solution on bark covering the entire circumference of the tree trunk. Wet the bark just to the point of saturation and run-off to the soil surface.

Certain surfactants may cause phytotoxicity upon direct contact with the foliage of non-target plants or the foliage on the tree that is being treated. Make sure the spray solution with surfactant does not contact foliage. Bark penetrating surfactants may change the color of lichens causing an objectionable appearance on the bark surface following treatment.

RESTRICTIONS

- Do not apply to wet bark, during rainfall, or if rain is expected within 12 hours after application.
- Do not make more than one application per year.
- Do not apply more than 0.77 lbs (0.54 lbs ai) Transtect 70WSP per acre per year.
- Application to heavily infested trees may not prevent the decline or death of trees due to existing damage and tree stress.
- Do not make bark applications when trees are dormant, under drought stress or not actively taking up water from the soil.

Applications should be made before trees are heavily infested and showing obvious symptoms of decline.

**Bark application for systemic insect control on ornamental plants in outdoor landscapes
(commercial, industrial, recreational and residential), parks, schools, cemeteries, airport grounds, military grounds,
golf courses, woodlots, athletic fields, forests (National, Private and State) and other wooded areas.**

Pests	Rate
Adelgids (such as: Hemlock Woolly) Flatheaded Borers (such as Emerald Ash, Appletree) Mountain Pine Beetle* Scale (armored and soft) (such as Calico, Cryptomeria, Elongate hemlock, Fig) Thrips (suppression) Whiteflies	<p>Bark Spray Use 6 – 12 (0.6 oz) packets (3.6 – 7.2 oz) of Transtect 70WSP per gallon of water. Consider using the higher rate for trees larger than 24 inches DBH.</p> <p>(Each packet contains approximately 0.026 lbs active ingredient)</p> <p>Spray bark on root flare (buttress roots) and on trunk between soil surface and 4 to 5 feet above the soil surface. Adjust nozzle to uniformly distribute spray over the entire circumference of the tree trunk and buttress roots. Wet bark just to the point of saturation and run off onto soil. Apply with a low volume sprayer operated at 10 to 20 PSI to prevent tree damage, bounce back and drift.</p> <p>Depending on bark type and thickness, 1 gal of spray solution will typically cover 65-85 inches of trunk diameter (1.5 – 2.0 fl oz/ inch of trunk diameter) when measured at 4.5 feet above the soil.</p> <p>For optimum control, apply to actively growing trees, and time application so that Transtect 70WSP has had time to move to insect feeding sites when target life stage is present.</p> <p>Control may be less effective in trees with thick bark and at times when trees are not actively growing or transpiring.</p> <p>*For Mountain Pine Beetle; apply between two weeks before and two weeks after peak of expected flight activity.</p>

Do not apply more than 0.77 lbs (0.54 lbs ai) per acre of landscape or forest per year.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

PESTICIDE STORAGE

Keep pesticide in original container.
 Do not put concentrate or dilute into food or drink containers.
 Store in a cool dry place.
 Do not store diluted spray.
 For help with any spill, leak, fire or exposure involving this material, call CHEMTREC
 day or night 1-800-424-9300.

PESTICIDE DISPOSAL

Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Nonrefillable outer bag. Do not reuse or refill the outer bag.
 Completely empty bag into application equipment. Offer for recycling, if available or dispose of empty bag in a sanitary landfill or by incineration.